

This listing of claims replaces all prior versions and listings of claims in this application.

**LISTING OF CLAIMS:**

---

Claims 1-12 (Canceled)

B2 Claim 13 (Original): A method for preparing a subcutaneously administrable biologically active agent, said method comprising:

(a) providing a biologically active agent which can exist in a native conformational state, a denatured conformational state, and an intermediate conformational state which is reversible to said native state and is conformationally between said native and denatured states; and

(b) exposing said biologically active agent to a complexing perturbant to reversibly transform said biologically active agent to said intermediate state and to form a subcutaneously administrable supramolecular complex,

said perturbant having a molecular weight ranging from about 150 to about 600 daltons, and having at least one hydrophilic moiety and at least one hydrophobic moiety,

said supramolecular complex comprising said perturbant non-covalently complexed with said biologically active agent; and

said biologically active agent not forming a microsphere with said perturbant;

wherein said perturbant is in an amount effective for subcutaneous

{M:\1946\1A483\00056406.DOC [REDACTED]}

Claim 14 (Original): A method as defined in claim 13, wherein said intermediate state has  $G$  ranging from about -20 kcal/mole to about 20 kcal/moles relative to said native state.

Claim 16 (Previously Presented): A method as defined in claim 15, wherein said biologically-active agent is selected from the group consisting of human growth hormone, bovine growth hormone, growth hormone-releasing hormone, an interferon, interleukin-II, insulin, heparin, calcitonin, erythropoietin, atrial natriuretic factor, an antigen, a monoclonal antibody, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, vasopressin, cromolyn sodium, vancomycin, desferrioxamine (DFO), low molecular weight heparin, antimicrobials, or any combination of any of the foregoing.

{M:\1946\1A483\00056406.DOC / 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 }


Claim 18 (Original): A method as defined in claim 13, wherein said perturbant is selected from the group consisting of an acylated amino acid and an acylated poly amino acid.

*B 2*  
*past*  
Claim 19 (Original): A method as defined in claim 13, wherein said perturbant is selected from the group consisting of a sulfonated amino acid and a sulfonated poly amino acid.

Claim 20 (Original): A method as defined in claim 13, wherein said perturbant is selected from the group consisting of an acylated aldehyde of an amino acid and an acylated aldehyde of a poly amino acid.

Claim 21 (Original): A method as defined in claim 13, wherein said perturbant is selected from the group consisting of an acylated ketone of an amino acid and an acylated ketone of a poly amino acid.

Claim 22 (Original): A method as defined in claim 13, wherein said perturbant comprises a carboxylic acid having the formula



wherein R is C<sub>1</sub> to C<sub>24</sub> alkyl, C<sub>2</sub> to C<sub>24</sub> alkenyl, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, phenyl, naphthyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)phenyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)phenyl, (C<sub>1</sub> to C<sub>10</sub> alkyl) naphthyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)naphthyl, phenyl(C<sub>1</sub> to C<sub>10</sub> alkyl), phenyl(C<sub>2</sub> to C<sub>10</sub> alkenyl), naphthyl(C<sub>1</sub> to C<sub>10</sub> alkyl) and naphthyl(C<sub>2</sub> to C<sub>10</sub> alkenyl);  
{M:\1946\1A483\00056406.DOC 1000000000000000 }

R being optionally substituted with C<sub>1</sub> to C<sub>10</sub> alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub> alkoxy, -OH, -SH, -CO<sub>2</sub>R<sup>1</sup>, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, heterocyclic having 3-10 ring atoms wherein the hetero atom is one or more atoms of N, O, S or any combination thereof, aryl, (C<sub>1</sub> to C<sub>10</sub> alkyl)aryl, aryl(C<sub>1</sub> to C<sub>10</sub>)alkyl, or any combination thereof;

*B1*  
*cat* R being optionally interrupted by oxygen, nitrogen, sulfur, or any combination thereof; and


R<sup>1</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>2</sub> to C<sub>4</sub> alkenyl; or  
a salt thereof.

Claim 23 (Original): A subcutaneous delivery composition comprising a supramolecular complex comprising:

(a) a biologically active agent in an intermediate conformational state non-covalently complexed with

(b) a complexing perturbant having a molecular weight ranging from about 150 to about 600 and having at least one hydrophilic moiety and at least one hydrophobic moiety;

wherein said intermediate state is reversible to said native state and is conformationally between a native conformational and a denatured conformational state of said biologically active agent and said composition is not a microsphere; and wherein said perturbant is in an amount effective for subcutaneous delivery of said biologically active agent.

{M:\1946\1A483\00056406.DOC  }

Claim 24 (Original): A composition as defined in claim 23, wherein said biologically active agent is selected from the group consisting of a peptide, a micropolysaccharide, a carbohydrate, a lipid, a pesticide, or any combination of the foregoing.

032  
rest  
Claim 25 (Previously Presented): A composition as defined in claim 24, wherein said biologically-active agent is selected from the group consisting of human growth hormone, bovine growth hormone, growth hormone-releasing hormone, an interferon, interleukin-II, insulin, heparin, calcitonin, erythropoietin, atrial naturetic factor, an antigen, a monoclonal antibody, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, vasopressin, cromolyn sodium, vancomycin, desferrioxamine (DFO), low molecular weight heparin, antimicrobials, or any combination of any of the foregoing.

Claim 26 (Original): A composition as defined in claim 23, wherein said perturbant comprises a proteinoid.

Claim 27 (Original): A composition as defined in claim 23, wherein said perturbant is selected from the group consisting of an acylated amino acid and an acylated poly amino acid.


Impure  
Claim 28 (Original): A method as defined in claim 46, wherein said perturbant is selected from the group consisting of a sulfonated amino acid and a {M:\1946\1A483\00056406.DOC [REDACTED] }

sulfonated poly amino acid.

Claim 29 (Original): A method as defined in claim 23, wherein said perturbant is selected from the group consisting of an acylated aldehyde of an amino acid and an acylated aldehyde of a poly amino acid.

B1  
cont  
Claim 30 (Original): A method as defined in claim 23, wherein said perturbant is selected from the group consisting of an acylated ketone of an amino acid and an acylated ketone of a poly amino acid.

Claim 31 (Original): A method as defined in claim 23, wherein said perturbant comprises a carboxylic acid having the formula



wherein R is C<sub>1</sub> to C<sub>24</sub> alkyl, C<sub>2</sub> to C<sub>24</sub> alkenyl, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, phenyl, naphthyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)phenyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)phenyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)naphthyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)naphthyl, phenyl(C<sub>1</sub> to C<sub>10</sub> alkyl), phenyl(C<sub>2</sub> to C<sub>10</sub> alkenyl), naphthyl(C<sub>1</sub> to C<sub>10</sub> alkyl) and naphthyl(C<sub>2</sub> to C<sub>10</sub> alkenyl);

R being optionally substituted with C<sub>1</sub> to C<sub>10</sub> alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub> alkoxy, -OH, -SH, -CO<sub>2</sub>R<sup>1</sup>, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, heterocyclic having 3-10 ring atoms wherein the hetero atom is one or more atoms of N, O, S or any combination thereof, aryl, (C<sub>1</sub> to C<sub>10</sub> alkyl)aryl, aryl(C<sub>1</sub> to C<sub>10</sub>)alkyl, or any combination thereof;

R being optionally interrupted by oxygen, nitrogen, sulfur, or any combination  
{M:\1946\1A483\00056406.DOC [REDACTED]}

thereof; and


R<sup>1</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>2</sub> to C<sub>4</sub> alkenyl; or  
a salt thereof.

Claim 32 (Original): A dosage unit form comprising:

- B.1*  
*Cont*
- (A) a composition as defined in claim 23; and
  - (B)
    - (a) an excipient,
    - (b) a diluent,
    - (c) a disintegrant,
    - (d) a lubricant,
    - (e) a plasticizer,
    - (f) a colorant,
    - (g) a dosing vehicle, or
    - (h) any combination thereof.

Claim 33 (Original): A method for preparing an agent which is capable of being administered by the subcutaneous route to a subject in need of said agent, said method comprising:

- (a) providing a biologically active agent which can exist in a native conformational state, a denatured conformational state, and an intermediate conformational state which is reversible to said native state and is conformationally between said native and denatured states;

{M:\1946\1A483\00056406.DOC  }

(b) exposing said biologically active agent to a complexing perturbant to reversibly transform said biologically active agent to said intermediate state and to form a subcutaneously administrable supramolecular complex,

said perturbant having a molecular weight between about 150 and about 600 daltons, and having at least one hydrophilic moiety and one hydrophilic moiety,

said supramolecular complex comprising said perturbant non-covalently complexed with said biologically active agent, and

*B2*  
*cont* said biologically active agent not forming a microsphere with said perturbant;

wherein said perturbant is in an amount effective for subcutaneous delivery of said biologically active agent; and

(c) preparing a mimetic of said supramolecular complex.

Claim 34 (Original): A method as defined in claim 33, wherein said biologically active agent comprises a peptide and said mimetic comprises a peptide mimetic.

Claim 35 (Original): A method for preparing an agent which is capable of being administered by the subcutaneous route to a subject in need of said agent, said method comprising:

(a) providing a biologically active agent which can exist in a native conformational state, a denatured conformational state, and an intermediate which is

{M:\1946\1A483\00056406.DOC [REDACTED]}



reversible to said native state and is conformationally between said native and denatured states;

(b) exposing said biologically active agent to a perturbant to reversibly transform said biologically active agent to said intermediate state, wherein said perturbant is in an amount effective for subcutaneous delivery of said biologically active agent; and

B.I.  
est (c) preparing a mimetic of said intermediate state.

Claim 36 (Original): A method as defined in claim 35, wherein said perturbant comprises a pH changing agent, an ionic strength changing agent, or guanidine hydrochloride.

Claim 37 (Currently Amended): A subcutaneous delivery composition comprising a mimetic, the mimetic being a mimetic of ~~the~~ a subcutaneous delivery composition prepared by the method of claim 13.

Claims 38-49 (Canceled)

Claim 50 (Original): A method for preparing a sublingually administrable biologically active agent, said method comprising:

(a) providing a biologically active agent which can exist in a native conformational state, a denatured conformational state, and an intermediate conformational state which is reversible to said native state and is conformationally

{M:\1946\1A483\00056406.DOC 10010000000000000000 }

between said native and denatured states; and

(b) exposing said biologically active agent to a complexing perturbant to reversibility transform said biologically active agent to said intermediate state and to form a sublingually administrable supramolecular complex,

said perturbant having a molecular weight ranging from about 150 to about 600 daltons, and having at least one hydrophilic moiety and at least one hydrophobic moiety,

said supramolecular complex comprising said perturbant non-covalently complexed with said biologically active agent; and

said biologically active agent not forming a microsphere with said perturbant;

wherein said perturbant is in an amount effective for sublingual delivery of said biologically active agent.

Claim 51 (Original): A method as defined in claim 50, wherein said intermediate state has  $\Delta G$  ranging from about -20 kcal/mole to about 20 kcal/moles relative to said native state.

Claim 52 (Original): A method as defined in claim 50, wherein said biologically active agent is selected from the group consisting of a peptide, a micropolysaccharide, a carbohydrate, a lipid, a pesticide, or any combination of the foregoing.

Claim 53 (Previously Presented): A method as defined in claim 52,  
{M:\1946\1A483\00056406.DOC [REDACTED] }

wherein said biologically-active agent is selected from the group consisting of human growth hormone, bovine growth hormone, growth hormone-releasing hormone, an interferon, interleukin-II, insulin, heparin, calcitonin, erythropoietin, atrial naturetic factor, an antigen, a monoclonal antibody, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, vasopressin, cromolyn sodium, vancomycin, desferrioxamine (DFO), low molecular weight heparin, antimicrobials, or any combination of any of the foregoing.

Claim 54 (Original): A method as defined in claim 50, wherein said perturbant comprises a proteinoid.

*By cont.*  
Claim 55 (Original): A method as defined in claim 50, wherein said perturbant is selected from the group consisting of an acylated amino acid and an acylated poly amino acid.


Claim 56 (Original): A method as defined in claim 50, wherein said perturbant is selected from the group consisting of a sulfonated amino acid and a sulfonated poly amino acid.

Claim 57 (Original): A method as defined in claim 50, wherein said perturbant is selected from the group consisting of an acylated aldehyde of an amino acid and an acylated aldehyde of a poly amino acid.

{M:\1946\1A483\00056406.DOC 10000000000000000000 }

Claim 58 (Original): A method as defined in claim 50, wherein said perturbant is selected from the group consisting of an acylated ketone of an amino acid and an acylated ketone of a poly amino acid.

Claim 59 (Original): A method as defined in claim 50, wherein said perturbant comprises a carboxylic acid having the formula



wherein R is C<sub>1</sub> to C<sub>24</sub> alkyl, C<sub>2</sub> to C<sub>24</sub> alkenyl, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, phenyl, naphthyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)phenyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)phenyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)naphthyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)naphthyl, phenyl(C<sub>1</sub> to C<sub>10</sub> alkyl), phenyl(C<sub>2</sub> to C<sub>10</sub> alkenyl), naphthyl(C<sub>1</sub> to C<sub>10</sub> alkyl) and naphthyl(C<sub>2</sub> to C<sub>10</sub> alkenyl);

*By cont.*  
R being optionally substituted with C<sub>1</sub> to C<sub>10</sub> alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub> alkoxy, -OH, -SH, -CO<sub>2</sub>R<sup>1</sup>, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, heterocyclic having 3-10 ring atoms wherein the hetero atom is one or more atoms of N, O, S or any combination thereof, aryl, (C<sub>1</sub> to C<sub>10</sub> alkyl)aryl, aryl(C<sub>1</sub> to C<sub>10</sub>)alkyl, or any combination thereof;

R being optionally interrupted by oxygen, nitrogen, sulfur, or any combination thereof; and

R<sup>1</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>2</sub> to C<sub>4</sub> alkenyl; or a salt thereof.

Claim 60 (Original): A sublingual delivery composition comprising a

{M:\1946\1A483\00056406.DOC [REDACTED]}

supramolecular complex comprising:

(a) a biologically active agent in an intermediate conformational state non-covalently complexed with

(b) a complexing perturbant having a molecular weight ranging from about 150 to about 600 and having at least one hydrophilic moiety and at least one hydrophobic moiety;

wherein said intermediate state is reversible to said native state and is conformationally between a native conformational and a denatured conformational state of said biologically active agent and said composition is not a microsphere; and wherein said perturbant is in an amount effective for sublingual delivery of said biologically active agent.

Claim 61 (Original): A composition as defined in claim 60, wherein said biologically active agent is selected from the group consisting of a peptide, a micropolysaccharide, a carbohydrate, a lipid, a pesticide, or any combination of the foregoing.

Claim 62 (Previously Presented): A composition as defined in claim 61, wherein said biologically-active agent is selected from the group consisting of human growth hormone, bovine growth hormone, growth hormone-releasing hormone, an interferon, interleukin-II, insulin, heparin, calcitonin, erythropoietin, atrial naturetic factor, an antigen, a monoclonal antibody, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, vasopressin, cromolyn sodium, vancomycin,

{M:\1946\1A483\00056406.DOC 10000000000000000000 }

desferrioxamine (DFO), low molecular weight heparin, antimicrobials, or any combination of any of the foregoing.

Claim 63 (Original): A composition as defined in claim 60, wherein said perturbant comprises a proteinoid.

Claim 64 (Original): A composition as defined in claim 60, wherein said perturbant is selected from the group consisting of an acylated amino acid and an acylated poly amino acid.

*By cont.*  
Claim 65 (Original): A method as defined in claim 60, wherein said perturbant is selected from the group consisting of a sulfonated amino acid and a sulfonated poly amino acid.

Claim 66 (Original): A method as defined in claim 60, wherein said perturbant is selected from the group consisting of an acylated aldehyde of an amino acid and an acylated aldehyde of a poly amino acid.

Claim 67 (Original): A method as defined in claim 60, wherein said perturbant is selected from the group consisting of an acylated ketone of an amino acid and an acylated ketone of a poly amino acid.

Claim 68 (Original): A method as defined in claim 60, wherein said  
{M:\1946\1A483\00056406.DOC [REDACTED]}

perturbant comprises a carboxylic acid having the formula



wherein R is C<sub>1</sub> to C<sub>24</sub> alkyl, C<sub>2</sub> to C<sub>24</sub> alkenyl, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, phenyl, naphthyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)phenyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)phenyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)naphthyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)naphthyl, phenyl(C<sub>1</sub> to C<sub>10</sub> alkyl), phenyl(C<sub>2</sub> to C<sub>10</sub> alkenyl), naphthyl(C<sub>1</sub> to C<sub>10</sub> alkyl) and naphthyl(C<sub>2</sub> to C<sub>10</sub> alkenyl);

B1  
cont.

R being optionally substituted with C<sub>1</sub> to C<sub>10</sub> alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub> alkoxy, -OH, -SH, -CO<sub>2</sub>R<sup>1</sup>, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, heterocyclic having 3-10 ring atoms wherein the hetero atom is one or more atoms of N, O, S or any combination thereof, aryl, (C<sub>1</sub> to C<sub>10</sub> alkyl)aryl, aryl(C<sub>1</sub> to C<sub>10</sub>)alkyl, or any combination thereof;

R being optionally interrupted by oxygen, nitrogen, sulfur, or any combination thereof; and

R<sup>1</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>2</sub> to C<sub>4</sub> alkenyl; or  
a salt thereof.

Claim 69 (Original): A dosage unit form comprising:

- (A) a composition as defined in claim 60; and
- (B)
  - (a) an excipient,
  - (b) a diluent,
  - (c) a disintegrant,
  - (d) a lubricant,

- B!  
cont.

(a) providing a biologically active agent which can exist in a native conformational state, a denatured conformational state, and an intermediate conformational state which is reversible to said native state and is conformationally between said native and denatured states;

said perturbant having a molecular weight between about 150 and about 600 daltons, and having at least one hydrophilic moiety and one hydrophilic moiety,

said biologically active agent not forming a microsphere with said  
perturbant;

{M:\1946\1A483\00056406.DOC 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 }



(c) preparing a mimetic of said supramolecular complex.

Claim 72 (Original): A method for preparing an agent which is capable of being administered by the sublingual route to a subject in need of said agent, said method comprising:

(b) exposing said biologically active agent to a perturbant to reversibly transform said biologically active agent to said intermediate state, wherein said perturbant is in an amount effective for sublingual delivery of said biologically active agent; and

Claim 73 (Original): A method as defined in claim 72, wherein said perturbant comprises a pH changing agent, an ionic strength changing agent, or guanidine hydrochloride.

-18-

Claim 74 (Currently Amended): ~~An oral~~ A sublingual delivery composition comprising a mimetic, the mimetic being a mimetic of the oral a sublingual delivery composition prepared by the method of claim 50.

Claims 75-86 (Canceled)

Claim 87 (Original): A method for preparing an intranasally administrable biologically active agent, said method comprising:

*B1 cont.* (a) providing a biologically active agent which can exist in a native conformational state, a denatured conformational state, and an intermediate conformational state which is reversible to said native state and is conformationally between said native and denatured states; and

(b) exposing said biologically active agent to a complexing perturbant to reversibility transform said biologically active agent to said intermediate state and to form an intranasally administrable supramolecular complex,

said perturbant having a molecular weight ranging from about 150 to about 600 daltons, and having at least one hydrophilic moiety and at least one hydrophobic moiety,

said supramolecular complex comprising said perturbant non-covalently complexed with said biologically active agent; and

said biologically active agent not forming a microsphere with said perturbant;

wherein said perturbant is in an amount effective for intranasal delivery of said

{M:\1946\1A483\00056406.DOC {XXXXXXXXXXXXXXXXXXXX} }

biologically active agent.

Claim 88 (Original): A method as defined in claim 87, wherein said intermediate state has  $\Delta G$  ranging from about -20 kcal/mole to about 20 kcal/moles relative to said native state.

Claim 89 (Original): A method as defined in claim 87, wherein said biologically active agent is selected from the group consisting of a peptide, a micropolysaccharide, a carbohydrate, a lipid, a pesticide, or any combination of the foregoing.

*B1 Cont*  
Claim 90 (Previously Presented): A method as defined in claim 89, wherein said biologically-active agent is selected from the group consisting of human growth hormone, bovine growth hormone, growth hormone-releasing hormone, an interferon, interleukin-II, insulin, heparin, calcitonin, erythropoietin, atrial natriuretic factor, an antigen, a monoclonal antibody, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, vasopressin, cromolyn sodium, vancomycin, desferrioxamine (DFO), low molecular weight heparin, antimicrobials, or any combination of any of the foregoing.

Claim 91 (Original): A method as defined in claim 87, wherein said perturbant comprises a proteinoid.

{M:\1946\1A483\00056406.DOC 10000000000000000000 }


Claim 92 (Original): A method as defined in claim 87, wherein said perturbant is selected from the group consisting of an acylated amino acid and an acylated poly amino acid.

Claim 93 (Original): A method as defined in claim 87, wherein said perturbant is selected from the group consisting of a sulfonated amino acid and a sulfonated poly amino acid.

Claim 94 (Original): A method as defined in claim 87, wherein said perturbant is selected from the group consisting of an acylated aldehyde of an amino acid and an acylated aldehyde of a poly amino acid.

Claim 95 (Original): A method as defined in claim 87, wherein said perturbant is selected from the group consisting of an acylated ketone of an amino acid and an acylated ketone of a poly amino acid.

Claim 96 (Original): A method as defined in claim 87, wherein said perturbant comprises a carboxylic acid having the formula



wherein R is C<sub>1</sub> to C<sub>24</sub> alkyl, C<sub>2</sub> to C<sub>24</sub> alkenyl, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, phenyl, naphthyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)phenyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)phenyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)naphthyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)naphthyl, phenyl(C<sub>1</sub> to C<sub>10</sub> alkyl), phenyl(C<sub>2</sub> to C<sub>10</sub> alkenyl), naphthyl(C<sub>1</sub> to C<sub>10</sub> alkyl) and naphthyl(C<sub>2</sub> to C<sub>10</sub> alkenyl);

{M:\1946\1A483\00056406.DOC 10000000000000000000 }

R being optionally substituted with C<sub>1</sub> to C<sub>10</sub> alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub> alkoxy, -OH, -SH, -CO<sub>2</sub>R<sup>1</sup>, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, heterocyclic having 3-10 ring atoms wherein the hetero atom is one or more atoms of N, O, S or any combination thereof, aryl, (C<sub>1</sub> to C<sub>10</sub> alkyl)aryl, aryl(C<sub>1</sub> to C<sub>10</sub>)alkyl, or any combination thereof;

R being optionally interrupted by oxygen, nitrogen, sulfur, or any combination thereof; and

R<sup>1</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>2</sub> to C<sub>4</sub> alkenyl; or  
a salt thereof.

*B1 cont*  
Claim 97 (Original): An intranasal delivery composition comprising a supramolecular complex comprising:

(a) a biologically active agent in an intermediate conformational state non-covalently complexed with

(b) a complexing perturbant having a molecular weight ranging from about 150 to about 600 and having at least one hydrophilic moiety and at least one hydrophobic moiety;

wherein said intermediate state is reversible to said native state and is conformationally between a native conformational and a denatured conformational state of said biologically active agent and said composition is not a microsphere; and wherein said perturbant is in an amount effective for intranasal delivery of said biologically active agent.

{M:\1946\1A483\00056406.DOC [REDACTED]}

Claim 98 (Original): A composition as defined in claim 97, wherein said biologically active agent is selected from the group consisting of a peptide, a micropolysaccharide, a carbohydrate, a lipid, a pesticide, or any combination of the foregoing.

Claim 99 (Previously Presented): A composition as defined in claim 98, wherein said biologically-active agent is selected from the group consisting of human growth hormone, bovine growth hormone, growth hormone-releasing hormone, an interferon, interleukin-II, insulin, heparin, calcitonin, erythropoietin, atrial naturetic factor, an antigen, a monoclonal antibody, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, vasopressin, cromolyn sodium, vancomycin, desferrioxamine (DFO), low molecular weight heparin, antimicrobials, or any combination of any of the foregoing.

Claim 100 (Original): A composition as defined in claim 97, wherein said perturbant comprises a proteinoid.

Claim 101 (Original): A composition as defined in claim 97, wherein said perturbant is selected from the group consisting of an acylated amino acid and an acylated poly amino acid.


Claim 102 (Original): A method as defined in claim 97, wherein said perturbant is selected from the group consisting of a sulfonated amino acid and a sulfonated poly amino acid.

{M:\1946\1A483\00056406.DOC [REDACTED] }

Claim 103 (Original): A method as defined in claim 97, wherein said perturbant is selected from the group consisting of an acylated aldehyde of an amino acid and an acylated aldehyde of a poly amino acid.

Claim 104 (Original): A method as defined in claim 97, wherein said perturbant is selected from the group consisting of an acylated ketone of an amino acid and an acylated ketone of a poly amino acid.

Claim 105 (Original): A method as defined in claim 97, wherein said perturbant comprises a carboxylic acid having the formula



*B1 cont*  
wherein R is C<sub>1</sub> to C<sub>24</sub> alkyl, C<sub>2</sub> to C<sub>24</sub> alkenyl, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, phenyl, naphthyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)phenyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)phenyl, (C<sub>1</sub> to C<sub>10</sub> alkyl)naphthyl, (C<sub>2</sub> to C<sub>10</sub> alkenyl)naphthyl, phenyl(C<sub>1</sub> to C<sub>10</sub> alkyl), phenyl(C<sub>2</sub> to C<sub>10</sub> alkenyl), naphthyl(C<sub>1</sub> to C<sub>10</sub> alkyl) and naphthyl(C<sub>2</sub> to C<sub>10</sub> alkenyl);

R being optionally substituted with C<sub>1</sub> to C<sub>10</sub> alkyl, C<sub>2</sub> to C<sub>10</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub> alkoxy, -OH, -SH, -CO<sub>2</sub>R<sup>1</sup>, C<sub>3</sub> to C<sub>10</sub> cycloalkyl, C<sub>3</sub> to C<sub>10</sub> cycloalkenyl, heterocyclic having 3-10 ring atoms wherein the hetero atom is one or more atoms of N, O, S or any combination thereof, aryl, (C<sub>1</sub> to C<sub>10</sub> alkyl)aryl, aryl(C<sub>1</sub> to C<sub>10</sub>)alkyl, or any combination thereof;

R being optionally interrupted by oxygen, nitrogen, sulfur, or any combination thereof; and

{M:\1946\1A483\00056406.DOC 10000000000000000000 }

R<sup>1</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>2</sub> to C<sub>4</sub> alkenyl; or  
a salt thereof.

Claim 106 (Original): A dosage unit form comprising:

- (A) a composition as defined in claim 97; and
- (B) (a) an excipient,  
(b) a diluent,  
(c) a disintegrant,  
(d) a lubricant,  
(e) a plasticizer,  
(f) a colorant,  
(g) a dosing vehicle, or  
(h) any combination thereof.

B<sub>1</sub>  
cont.

Claim 107 (Original): A method for preparing an agent which is capable of  
being administered by the intranasal route to a subject in need of said agent, said  
method comprising:

- (a) providing a biologically active agent which can exist in a native  
conformational state, a denatured conformational state, and an intermediate  
conformational state which is reversible to said native state and is conformationally  
between said native and denatured states;
- (b) exposing said biologically active agent to a complexing perturbant to

{M:\1946\1A483\00056406.DOC 10000000000000000000 }



reversibly transform said biologically active agent to said intermediate state and to form an intranasally administrable supramolecular complex,

said perturbant having a molecular weight between about 150 and about 600 daltons, and having at least one hydrophilic moiety and one hydrophilic moiety,

said supramolecular complex comprising said perturbant non-covalently complexed with said biologically active agent, and

said biologically active agent not forming a microsphere with said perturbant;

wherein said perturbant is in an amount effective for intranasal delivery of said biologically active agent; and

(c) preparing a mimetic of said supramolecular complex.

B1  
cont.

Claim 108 (Original): A method as defined in claim 107, wherein said biologically active agent comprises a peptide and said mimetic comprises a peptide mimetic.

Claim 109 (Original): A method for preparing an agent which is capable of being administered by the intranasal route to a subject in need of said agent, said method comprising:

(a) providing a biologically active agent which can exist in a native conformational state, a denatured conformational state, and an intermediate which is reversible to said native state and is conformationally between said native and denatured states;

{M:\1946\1A483\00056406.DOC [REDACTED]}

(c) preparing a mimetic of said intermediate state.

Claim 111 (Original): An ~~oral~~ intranasal delivery composition comprising a mimetic, the mimetic being a mimetic of the oral ~~an~~ intranasal delivery composition prepared by the method of claim 87.

Claim 113 (Currently Amended): The method of claim 55 128, wherein the biologically active agent is growth-hormone releasing hormone.

Claim 114 (Currently Amended): The method of claim ~~55~~ 128, wherein the biologically active agent is insulin.

Claim 115 (Currently Amended): The method of claim ~~55~~ 128, wherein the biologically active agent is heparin.

Claim 116 (Currently Amended): The method of claim ~~55~~ 128, wherein the biologically active agent is low molecular weight heparin.

Claim 117 (Currently Amended): The method of claim ~~55~~ 128, wherein the biologically active agent is calcitonin.


Claim 118 (Currently Amended): The method of claim ~~55~~ 128, wherein the biologically active agent is cromolyn sodium.

*B!  
cont.*  
Claim 119 (Currently Amended): The method of claim ~~55~~ 128, wherein the biologically active agent is an antimicrobial.

Claim 120 (Currently Amended): The composition of claim ~~64~~ 129, wherein the biologically active agent is human growth hormone.

Claim 121 (Currently Amended): The composition of claim ~~64~~ 129, wherein the biologically active agent is growth-hormone releasing hormone.

Claim 122 (Currently Amended): The composition of claim ~~64~~ 129, wherein the biologically active agent is insulin.

{M:\1946\1A483\00056406.DOC  }

Claim 123 (Currently Amended): The composition of claim 64 129, wherein the biologically active agent is heparin.

Claim 124 (Currently Amended): The composition of claim 64 129, wherein the biologically active agent is low molecular weight heparin.

Claim 125 (Currently Amended): The composition of claim 64 129, wherein the biologically active agent is calcitonin.


Claim 126 (Currently Amended): The composition of claim 64 129, wherein the biologically active agent is cromolyn sodium.

*B1 cont.*  
Claim 127 (Currently Amended): The composition of claim 64 129, wherein the biologically active agent is an antimicrobial.

Claim 128 (New): A method as defined in claim 55, wherein said perturbant is an acylated amino acid.

Claim 129 (New): A composition as defined in claim 64, wherein said perturbant is an acylated amino acid.

Claim 130 (New): A method as defined in claim 128, wherein the biologically active agent is a peptide.

{M:\1946\1A483\00056406.DOC  }

Claim 131 (New): A method as defined in claim 130, wherein the biologically active agent is an interferon.

Claim 132 (New): A method as defined in claim 130, wherein the biologically active agent is erythropoietin.

Claim 133 (New): A method as defined in claim 130, wherein the biologically active agent is an antigen.

Claim 134 (New): A composition as defined in claim 129, wherein the biologically active agent is a peptide.

*B1 cont.*  
Claim 135 (New): A composition as defined in claim 134, wherein the biologically active agent is an interferon.

Claim 136 (New): A composition as defined in claim 134, wherein the biologically active agent is erythropoietin.

Claim 137 (New): A composition as defined in claim 134, wherein the biologically active agent is an antigen.

Claim 138 (New): A method as defined in claim 18, wherein said perturbant is an acylated amino acid.

{M:\1946\1A483\00056406.DOC [REDACTED] }

Claim 139 (New): The method of claim 138, wherein the biologically active agent is human growth hormone.

Claim 140 (New): The method of claim 138, wherein the biologically active agent is growth-hormone releasing hormone.

Claim 141 (New): The method of claim 138, wherein the biologically active agent is insulin.

Claim 142 (New): The method of claim 138, wherein the biologically active agent is heparin.

Claim 143 (New): The method of claim 138, wherein the biologically active agent is low molecular weight heparin.

Claim 144 (New): The method of claim 138, wherein the biologically active agent is calcitonin.

Claim 145 (New): The method of claim 138, wherein the biologically active agent is cromolyn sodium.

Claim 146 (New): The method of claim 138, wherein the biologically active agent is an antimicrobial.

{M:\1946\1A483\00056406.DOC [REDACTED]}

Claim 147 (New): A method as defined in claim 138, wherein the biologically active agent is a peptide.

Claim 148 (New): A method as defined in claim 147, wherein the biologically active agent is an interferon.

Claim 149 (New): A method as defined in claim 147, wherein the biologically active agent is erythropoietin.

Claim 150 (New): A method as defined in claim 147, wherein the biologically active agent is an antigen.

*B1 cont.*  
Claim 151 (New): A composition as defined in claim 27, wherein said perturbant is an acylated amino acid.

Claim 152 (New): The composition of claim 151, wherein the biologically active agent is human growth hormone.

Claim 153 (New): The composition of claim 151, wherein the biologically active agent is growth-hormone releasing hormone.

Claim 154 (New): The composition of claim 151, wherein the biologically active agent is insulin.

{M:\1946\1A483\00056406.DOC [REDACTED] }

Claim 155 (New): The composition of claim 151, wherein the biologically active agent is heparin.

Claim 156 (New): The composition of claim 151, wherein the biologically active agent is low molecular weight heparin.

Claim 157 (New): The composition of claim 151, wherein the biologically active agent is calcitonin.

*B1 cont.*  
Claim 158 (New): The composition of claim 151, wherein the biologically active agent is cromolyn sodium.

Claim 159 (New): The composition of claim 151, wherein the biologically active agent is an antimicrobial.

Claim 160 (New): A composition as defined in claim 151, wherein the biologically active agent is a peptide.

Claim 161 (New): A composition as defined in claim 160, wherein the biologically active agent is an interferon.

Claim 162 (New): A composition as defined in claim 160, wherein the biologically active agent is erythropoietin.

{M:\1946\1A483\00056406.DOC [REDACTED]}



Claim 163 (New): A composition as defined in claim 160, wherein the biologically active agent is an antigen.

Claim 164 (New): A method as defined in claim 92, wherein said perturbant is an acylated amino acid.

Claim 165 (New): The method of claim 164, wherein the biologically active agent is human growth hormone.

Claim 166 (New): The method of claim 164, wherein the biologically active agent is growth-hormone releasing hormone.

Claim 167 (New): The method of claim 164, wherein the biologically active agent is insulin.

Claim 168 (New): The method of claim 164, wherein the biologically active agent is heparin.

Claim 169 (New): The method of claim 164, wherein the biologically active agent is low molecular weight heparin.

Claim 170 (New): The method of claim 164, wherein the biologically active agent is calcitonin.

{M:\1946\1A483\00056406.DOC [REDACTED]}



Claim 179 (New): The composition of claim 177, wherein the biologically active agent is growth-hormone releasing hormone.

Claim 180 (New): The composition of claim 177, wherein the biologically active agent is insulin.

Claim 181 (New): The composition of claim 177, wherein the biologically active agent is heparin.


*B1  
cont* Claim 182 (New): The composition of claim 177, wherein the biologically active agent is low molecular weight heparin.

Claim 183 (New): The composition of claim 177, wherein the biologically active agent is calcitonin.

Claim 184 (New): The composition of claim 177, wherein the biologically active agent is cromolyn sodium.

Claim 185 (New): The composition of claim 177, wherein the biologically active agent is an antimicrobial.

Claim 186 (New): A composition as defined in claim 177, wherein the biologically active agent is a peptide.

{M:\1946\1A483\00056406.DOC  }

Claim 188 (New): A composition as defined in claim 186, wherein the biologically active agent is erythropoietin.

Claim 189 (New): A composition as defined in claim 186, wherein the biologically active agent is an antigen.